

10656
HAGIOXENUS MAYRI comb. nov.

By William L. Brown, Jr.
Department of Entomology
Cornell University
Ithaca, New York, USA

Pilot Register of Zoology
Card No. 19
Issued 20 May 1964.

W.L. Brown, Jr.
COLLECTION

pro Rhoptrymex Mayri Forel, 1912, Zool. Jahrb., suppl. 15, 1: 57, alate female. Type locality Poona, India.
Type in Muséum d'Histoire Naturelle, Geneva, examined 1963.

The R. mayri type is an aberrant form, obviously a parasite of some kind, but it is much more similar to the types of Hagioxenus schmitzi than to any of the known Rhoptrymex females; furthermore, as Forel noted in the original description, the radial cell is closed.

From H. schmitzi, H. mayri is distinguished by its much more abundant pilosity and by other relatively small differences. The type was collected "together with Pheidole latinoda."

Parasitic myrmecines are frequently much modified, with loss or reduction of some characters, and are particularly likely to converge in habitus. Even the rather close resemblance of H. schmitzi and H. mayri could of course be due to convergence.

Published with the aid of a grant from the Department of Entomology and Limnology, New York State College of Agriculture at Cornell University, Ithaca, New York. Edited by W. L. Brown, Jr.

A89890X
R1
R2
R3
R4
R5
R6
R7
R8
R9
R10
R11
R12
R13
R14
R15
R16
R17
R18
R19
R20
R21
R22
R23
R24
R25
R26
R27
R28
R29